

Nagoya University  
G30 Preparatory Lecture  
**Mathematics**

**Course I : Functions and Equations**

**Course II : Calculus**

**Course III: Linear Algebra**

# Course I : Functions and Equations

# Flow of This Lecture Video

1. **Lecture A** (About 20 min.)
  - Listen to a lecture.
2. Exercise A
  - Practice solving related problems.
  - Pause the video and solve the problem **by yourself**.
  - If you cannot, see the preceding lecture repeatedly.
3. Explanation of the answers (About 5 min.)
4. **Lecture B** (About 20 min.)
5. Exercise B
6. Explanation of the answers

# Contents

<b>Lesson 01</b>	Polynomials and Factoring
<b>Lesson 02</b>	Algebraic Equations and Functions
<b>Lesson 03</b>	Linear and Quadratic Inequalities
<b>Lesson 04</b>	Trigonometric Functions (I)
<b>Lesson 05</b>	Trigonometric Functions (II)
<b>Lesson 06</b>	Trigonometric Functions (III)
<b>Lesson 07</b>	Complex Numbers
<b>Lesson 08</b>	Exponential Functions
<b>Lesson 09</b>	Fractional Functions and Irrational Functions
<b>Lesson 10</b>	Inverse Functions
<b>Lesson 11</b>	Common Logarithms
<b>Lesson 12</b>	Law of Sines and Law of Cosines
<b>Lesson 13</b>	Applications of Trigonometric Functions
<b>Lesson 14</b>	Graphs and Equations (I)
<b>Lesson 15</b>	Graphs and Equations (II)

# Lesson 1

## Polynomials and Factoring


### 1A

- Polynomials
- Addition, Subtraction, and Multiplication

# Polynomials

**Q.** What is the area of a circle with radius  $r$  cm ?

**A.**  $\pi r^2$  (cm<sup>2</sup>) : Monomial




**Q.** What is the price for 5 shortcakes of  $x$  yen and 2 cream puffs of  $y$  yen?

**A.**  $5x + 2y$  (yen) : Polynomial



**Polynomial** is an expression of finite length constructed from variables and constant, using only the operation of addition, subtraction and multiplication.

## Some terminologies



# Some Notes

**Note 1 :** Degree of a polynomial is the largest degree of any one term.  
(Ex: Degree of  $2x^3 + 2x + 1$  is 3.)

**Note 2 :** The following terms are **not** polynomials.

$$3x^{\frac{1}{2}} + 1 \quad (= 3\sqrt{x} + 1) \qquad 2x^{-2} \quad (= 1/x^2)$$

**Note 3 :** Simplify by combining **like terms**.

$$5x^3 + 2x + 1 - 2x^2 + x \quad \rightarrow \quad 3x^3 - 2x^2 + 3x + 1$$

**Note 4 :** Write in the descending order of the degree

$$2x^3 + 2x + 1 - x^2 \quad \rightarrow \quad 2x^3 - x^2 + 2x + 1$$



# Basic Process of Multiplication

(two-term)×(two-term)

$$\begin{aligned}
 (ax + b)(cx + d) &= acx^2 + adx + bcx + bd \\
 &= acx^2 + (ad + bc)x + bd
 \end{aligned}$$

Steps ↓

- ① First terms
- ② Outer terms
- ③ Inner terms
- ④ Last terms

In general

Use the distribution property and combine.

**Example 1.** Find the product  $(2x^2 + x - 3)(x + 2)$

**Ans.**

<Method 1>

$$\begin{aligned}
 &(2x^2 + x - 3)(x + 2) \\
 &= (2x^2 + x - 3)x + (2x^2 + x - 3) \cdot 2 \\
 &= 2x^3 + x^2 - 3x + 4x^2 + 2x - 6 \\
 &= 2x^3 + 5x^2 - x - 6
 \end{aligned}$$

<Method 2>

$$\begin{array}{r}
 2x^2 + x - 3 \\
 \times) \quad x + 2 \\
 \hline
 2x^3 + x^2 - 3x \\
 4x^2 + 2x - 6 \\
 \hline
 2x^3 + 5x^2 - x - 6
 \end{array}$$



# Some Formulas

Memorize the following formulas for quick calculation.

$$\begin{array}{l} 1. \quad (a + b)^2 = a^2 + 2ab + b^2 \\ \quad \quad (a - b)^2 = a^2 - 2ab + b^2 \end{array} \quad \begin{array}{c} \boxed{\phantom{00}} \\ \leftarrow \end{array} \quad b \rightarrow (-b)$$

$$2. \quad (a + b)(a - b) = a^2 - b^2$$

$$3. \quad (x + a)(x + b) = x^2 + (a + b)x + ab$$

$$4. \quad (ax + b)(cx + d) = acx^2 + (ad + bc)x + bd$$

$$\begin{array}{l} 5. \quad (a + b)^3 = a^3 + 3a^2b + 3ab^2 + b^3 \\ \quad \quad (a - b)^3 = a^3 - 3a^2b + 3ab^2 - b^3 \end{array} \quad \begin{array}{c} \boxed{\phantom{00}} \\ \leftarrow \end{array} \quad b \rightarrow (-b)$$

# Exercise

**Exercise 1.** Perform the indicated operation for the following polynomials.

$$A = 2x^3 - x^2 - 1, \quad B = -x^2 + 2x + 1$$

- (1) Add  $A$  and  $B$ .
- (2) Subtract  $B$  from  $A$ .
- (3) Multiply  $A$  and  $B$ .

Pause the video and solve the problem by yourself.

# Answer to the Exercise

**Exercise 1.** Perform the indicated operation for the following polynomials.

$$A = 2x^3 - x^2 - 1, \quad B = -x^2 + 2x + 1$$

(1) Add  $A$  and  $B$  .      (2) Subtract  $B$  from  $A$  .      (3) Multiply  $A$  and  $B$  .

**Ans.**

$$\begin{array}{r} (1) \quad 2x^3 - x^2 \quad -1 \\ +) \quad -x^2 + 2x + 1 \\ \hline 2x^3 - 2x^2 + 2x \end{array}$$

$$\begin{array}{r} (2) \quad 2x^3 - x^2 \quad -1 \\ -) \quad -x^2 + 2x + 1 \\ \hline 2x^3 \quad -2x - 2 \end{array}$$

$$\begin{array}{r} (3) \quad 2x^3 - x^2 \quad -1 \\ \times) \quad -x^2 + 2x + 1 \\ \hline -2x^5 + x^4 \quad + x^2 \\ + 4x^4 - 2x^3 \quad - 2x \\ + 2x^3 - x^2 \quad - 1 \\ \hline -2x^5 + 5x^4 \quad - 2x - 1 \end{array}$$

# Lesson 1

## Polynomials and Factoring

### 1B

- Factoring Polynomials
- How to factor polynomials
- Factor theorem

# What is Factoring ?

Factoring a polynomial is the opposite process of multiplying polynomials.

Expansion



$$(x + 2)(x + 3) = x^2 + 5x + 6$$



Factoring



## Factoring a number

$$12 = 2 \times 6$$

$$12 = \frac{1}{2} \times 24$$

Prime numbers

$$12 = \underline{2 \times 2 \times 3}$$

Completely factored

## Factoring a polynomial

$$x^4 - 16 = (x^2 + 4)(x^2 - 4) = \underline{(x^2 + 4)(x + 2)(x - 2)}$$

Completely factored

# How to Factor Polynomials

There is no perfect method to succeed factoring. Sometimes **insight** will help. However, there are several useful rules as follows.

## 1. Factor out the largest common factor.

$$[\text{Ex.}] \quad 2x^3 + 10x^2 + 12x = 2x(x^2 + 5x + 6) \quad (a + b) \quad \leftarrow \text{Rule 1}$$

## 2. Use a formula of factoring.

$$\begin{aligned} [\text{Ex.}] \quad 2x^3 + 10x^2 + 12x &= 2x(x^2 + 5x + 6) \\ &= 2x(x + 2)(x + 3) \end{aligned} \quad \leftarrow \text{Rule 2}$$

*Diagram annotations:* Red boxes labeled  $a$  and  $b$  point to the terms  $2x$  and  $6$  in the second line. Red boxes labeled  $ab$  and  $(a + b)$  point to the terms  $5x$  and  $6$  in the first line.

## 3. Rearrange by the variable with the lowest degree.

$$\begin{aligned} [\text{Ex.}] \quad x^3 + 3x^2 + 2xy + 6y &= (2x + 6)y + (x^3 + 3x^2) \quad \leftarrow \text{Rule 3} \\ &= 2y(x + 3) + x^2(x + 3) \quad \leftarrow \text{Rule 1} \\ &= (x^2 + 2y)(x + 3) \quad \leftarrow \text{Rule 1} \end{aligned}$$

## 4. Use “Factor Theorem”

To be discussed later.

# Several Advanced Examples

**Example 2.** Factor the following polynomials.

(1)  $2x^2 + 5xy + 3y^2 - 3x - 5y - 2$

(2)  $x^4 - 10x^2 + 9$

**Ans.**

(1) Every valuable has the same degree. In such a case, rearrange by any one of them.

$$2x^2 + 5xy + 3y^2 - 3x - 5y - 2$$

$$= 2x^2 + (5y - 3)x + (3y^2 - 5y - 2)$$

$$= 2x^2 + (5y - 3)x + (y - 2)(3y + 1)$$

$$= \{x + (y - 2)\}\{2x + (3y + 1)\}$$

$$= (x + y - 2)(2x + 3y + 1)$$

$$\begin{array}{rcl} 1 & \times & y - 2 \rightarrow 2y - 4 \\ 2 & \times & 3y + 1 \rightarrow 3y + 1 \\ \hline & & 5y - 3 \end{array}$$

(2) The polynomial has 4<sup>th</sup> degree. In such a case, observe the form carefully and use the similarity to the polynomial with a lower degree.

Put  $X = x^2$

Then  $x^4 - 10x^2 + 9 = X^2 - 10X + 9 = (X - 1)(X - 9)$

$$= (x^2 - 1)(x^2 - 9) = (x + 1)(x - 1)(x + 3)(x - 3)$$

# Factor Theorem

## Factor Theorem

A polynomial  $f(x)$  has a factor  $(x - k)$  if and only if  $f(k) = 0$ .

This theorem is commonly applied to the problems of **factoring a polynomial** and **finding the roots of a polynomial equation** (this will be explained later.)

### Steps of application:

1. **Guess** a number  $k$  and confirm  $f(k) = 0$ .
2. Divide  $f(x)$  by  $(x - k)$  and obtain  $g(x) = f(x)/(x - k)$ .
3. Then,  $f(x)$  is factored to  $f(x) = (x - k)g(x)$ .
4. It is easier to find factors of  $g(x)$  than that of  $f(x)$ .

**Example 3.** Factor the polynomial  $f(x) = x^3 + 4x^2 + x - 6$ .

**Ans.**  $f(1) = 0$  Therefore

$$\begin{aligned}x^3 + 4x^2 + x - 6 &= (x - 1)(x^2 + 5x + 6) \\ &= (x - 1)(x - 2)(x - 3)\end{aligned}$$

$$\begin{array}{r}x^2 + 5x + 6 \\ x - 1 \overline{) x^3 + 4x^2 + x - 6} \\ \underline{x^3 - x^2} \phantom{+ x - 6} \\ 5x^2 + x - 6 \\ \underline{5x^2 - 5x} \phantom{- 6} \\ 6x - 6 \\ \underline{6x - 6} \\ 0\end{array}$$



# Exercise

**Exercise 2.** Factor the following polynomials.

(1)  $4x^2 + 8x - 21$       (2)  $x^3 + 7x^2 + 8x + 2$

Pause the video and solve the problem by yourself.

# Answer to the Exercise

**Exercise 2.** Factor the following polynomials.

(1)  $4x^2 + 8x - 21$       (2)  $x^3 + 7x^2 + 8x + 2$

**Ans.**

- (1) Remember the formula  $(ax + b)(cx + d) = acx^2 + (ad + bc)x + bd$   
By trials, we find that  $a = 2$ ,  $b = -3$ ,  $c = 2$ ,  $d = 7$  satisfy the condition.  
(Refer to the right-side calculation)

Therefore  $4x^2 + 8x - 21 = (2x - 3)(2x + 7)$

$$\begin{array}{rcl} 2 & \times & -3 \rightarrow -6 \\ 2 & \times & 7 \rightarrow 14 \\ \hline 4 & -21 & 8 \end{array}$$

- (2) We use Factor Theorem.

By trials, we find  $x = -1$  satisfy  $f(-1) = 0$ .

By dividing  $x^3 + 7x^2 + 8x + 2$  by  $x + 1$ , we have

$$\frac{x^3 + 7x^2 + 8x + 2}{x + 1} = x^2 + 6x + 2$$

Therefore

$$x^3 + 7x^2 + 8x + 2 = (x^2 + 6x + 2)(x + 1)$$