

Lesson 9

Antiderivatives

9A

- Antiderivative

Inverse Problem

Inverse Problem

To find the function when its derivative is given.

$$[\text{Ex.}] \text{ velocity } v(t) \left(= \frac{ds(t)}{dt} \right) \rightarrow \text{ position } s(t)$$

Antiderivative

A function $F(x)$ is an **antiderivative** (or **primitive integral**) of $f(x)$ if $F'(x) = f(x)$.

[Ex.] $F(x) = x^3$ is an antiderivative of $f(x) = 3x^2$ because

$$\frac{d}{dx} (x^3) = 3x^2$$

Indefinite Integral

Example:

x^3 , $x^3 - 2$, $x^3 + 2$: antiderivatives of $3x^2$

Theorem

When $F'(x) = f(x)$, every other antiderivative is of the form


$$F(x) + C \quad (C : \text{constant})$$

Indefinite Integral

The collection of all antiderivatives of a function is called the **general antiderivative** or **indefinite integral** of $f(x)$ and denoted by $\int f(x)dx$

Namely,

$$\int f(x)dx = F(x) + C$$



Fundamental Indefinite Integrals

$$x^3 + C \begin{array}{c} \xrightarrow{\text{Differentiation}} \\ \xleftarrow{\text{Integration}} \end{array} 3x^2$$

Algebraic Functions

< Differentiation >

$$\frac{d}{dx} x^{n+1} = (n+1)x^n$$



< Integration >

$$\int x^n dx = \frac{x^{n+1}}{n+1} + C \quad \text{for } n \neq -1$$

Example

$$\int x^5 dx = \frac{1}{6} x^6 + C$$

Note

$$\text{For } n = -1 \quad \int x^{-1} dx = \frac{x^{-1+1}}{-1+1} + C = \frac{x^0}{0} + C$$

Meaningless
4

Graphical Interpretation.

Differentiation

Graph :
 $y = f(x)$



Slope :
 $\frac{df(x)}{dx}$


Integration

Slope :

$\frac{df(x)}{dx}$



Graph :
 $y = f(x)$




<Another interpretation>

Graph : $y = f(x)$  Area


(We will study this later)

Graphical Interpretation of Differentiation

How to find **the derivative** $y = \frac{1}{x}$ graphically from $y = \ln x$






$$y = \ln x$$



$$y = \frac{d}{dx} \ln x = \frac{1}{x}$$

Graphical Interpretation of Integration

How to find **the indefinite function** graphically from $y = \frac{1}{x}$



$$y = \int \frac{1}{x} dx$$

Algebraic derivation

For $x > 0$ (Defined domain of $y = \ln x$)

$$\frac{d}{dx} \ln x = \frac{1}{x} \quad \longleftrightarrow \quad \int \frac{1}{x} dx = \ln x + C$$

For $x < 0$

From the graphical discussion, we assume $y = \ln|x|$

Put $x = -u$ ($u > 0$)

From the chain rule

$$\frac{dy}{dx} = \frac{d(\ln|x|)}{dx} = \frac{d(\ln u)}{du} \frac{du}{dx} = \frac{1}{u} \cdot (-1) = -\frac{1}{u} = \frac{1}{x}$$

For all $x \neq 0$ $\int \frac{1}{x} dx = \ln|x| + C$

Trigonometric Integrals

$$\frac{d}{dx} \sin x = \cos x \quad \Rightarrow \quad \int \cos x dx = \sin x + C$$

$$\frac{d}{dx} \cos x = -\sin x \quad \Rightarrow \quad \int \sin x dx = -\cos x + C$$

$$\frac{d}{dx} \tan x = \frac{1}{\cos^2 x} \quad \Rightarrow \quad \int \frac{1}{\cos^2 x} dx = \tan x + C$$

Exponential Integrals

$$\frac{d}{dx} e^x = e^x \quad \rightarrow \quad \int e^x dx = e^x + C$$

$$\frac{d}{dx} a^x = a^x \ln a \quad \rightarrow \quad \int a^x dx = \frac{a^x}{\ln a} + C$$

Basic Rules of Integration

Sum Rule

$$\int \{f(x) + g(x)\} dx = \int f(x) dx + \int g(x) dx$$

Constant Multiple Rule

$$\int cf(x) dx = c \int f(x) dx$$

Example

Examples 9-2 Find the indefinite integrals of the following function.

(1) $(x+1)(x-2)$

(2) $\sin x + 2 \cos x$

Ans.

$$\begin{aligned} (1) \quad \int (x+1)(x-2) dx &= \int (x^2 - x - 2) dx = \int x^2 dx - \int x dx - 2 \int dx \\ &= \frac{x^3}{3} - \frac{x^2}{2} - 2x + C \end{aligned}$$

$$\begin{aligned} (2) \quad \int (\sin x + 2 \cos x) dx &= \int \sin x + 2 \int \cos x dx \\ &= -\cos x + 2 \sin x + C \end{aligned}$$

Exercise

Exercise. 9-1 The slope of the curve $y=f(x)$ is given by $3x^2$ and this curve passes the point $(1,2)$. Determine the equation of this curve.

Ans.

Pause the video and solve by yourself.

Answer to the Exercises

Exercise 12-1] The slope of the curve $y=f(x)$ is given by $3x^2$ and this curve passes the point $(1,2)$. Determine the equation of this curve.

Ans.


$$f'(x)=3x^2$$

$$\therefore f(x)=\int 3x^2 dx = x^3 + C$$

Since this line passes the point $(1,2)$,

$$f(1)=1+C=2 \quad \therefore C=1$$

$$\therefore y=x^3+1$$



Lesson 9

Antiderivatives

9B

- Substitution Method
- Integration by Parts (IBP)

Substitution Method (1)

Applicable when the integrand is of the form $f(u(x))u'(x)$

Example:

$(x^2 + 1)^5 x$	$\sin^5 x \cos x$
$u = x^2 + 1$	$u = \sin x$
$u' = 2x$	$u' = \cos x$

Let $F'(u) = f(u)$ or $F(u) = \int f(u)du$ ①

From the Chain Rule $\frac{d}{dx} F(u(x)) = F'(u)u'(x) = f(u)u'(x)$

By integration, we have $F(u(x)) = \int f(u)u'(x)dx$ ②

From ① and ②, we have

Substitution Method

$$\int f(u(x))u'(x)dx = \int f(u)du$$

Substitution Using Differentials

Differential

$$u = u(x) \longrightarrow \frac{du}{dx} = u'(x) \longrightarrow \textcircled{du} = u'(x)\textcircled{dx}$$
$$du = \frac{du}{\cancel{dx}} \cancel{dx}$$

< Example > If $u = x^3$, then $du = 3x^2 dx$

$$\int f(u) \underline{u'(x) dx} = \int f(u) \underline{du}$$

$du = \frac{du}{dx} dx$

← →

Example

Examples 9-4 Find the indefinite integral of the following functions.

$$(1) \int (x^2 + 1)^5 x dx \quad (2) \int \sin^5 x \cos x dx$$

Ans. (1) We put $x^2 + 1 = u$

$$\text{Then } 2x \frac{dx}{du} = 1 \quad \therefore x dx = \frac{1}{2} du$$

Substitution

$$\int (x^2 + 1)^5 x dx = \int u^5 \frac{1}{2} du = \frac{1}{12} u^6 + C = \frac{1}{12} (x^2 + 1)^6 + C$$

(2) We put $\sin x = u$

$$\text{Then } \cos x \frac{dx}{du} = 1 \quad \therefore \cos x dx = \frac{1}{\cos x} du$$

Substitution

$$\int \sin^5 x \cos x dx = \int u^5 du = \frac{1}{6} u^6 + C = \frac{1}{6} \sin^6 x + C$$

Substitution Method (2)

Substitution Method

$$\int f(x)dx = \int f(x(t))x'(t)dt$$

This is also written as follows.

$$\int f(x)dx = \int f(x(t))\frac{dx}{dt}dt$$

Example

Examples 9-5 Find the indefinite integral of the following functions.

$$\int x\sqrt{1-x}dx$$

Ans. We put $\sqrt{1-x} = t$. Then $x = 1 - t^2$ and $\frac{dx}{dt} = -2t$.

Therefore

$$\int x\sqrt{1-x}dx = \int (1-t^2)t(-2t)dt = \int (1-t^2)t(-2t)dt$$

$$= 2\int (t^4 - t^2)dt = 2\left(\frac{t^5}{5} - \frac{t^3}{3}\right) + C = -\frac{2}{15}(2+3x)(1-x)\sqrt{1-x} + C$$

Integration by Parts

We studied the derivative of product

$$\{f(x)g(x)\}' = f'(x)g(x) + f(x)g'(x)$$

After rearrangement, we have

$$f(x)g'(x) = \{f(x)g(x)\}' - f'(x)g(x)$$

Therefore, we obtain the following formula

Integration by Parts

$$\int f(x)g'(x)dx = f(x)g(x) - \int f'(x)g(x)dx$$

Example

Examples 9-5 Evaluate the following Integral.

$$\int x \cos x dx$$

Ans.

$$\int \overbrace{x}^{\boxed{f}} \overbrace{\cos x}^{\boxed{g'}} dx = \overbrace{x}^{\boxed{f}} \overbrace{\sin x}^{\boxed{g}} - \int \overbrace{1}^{\boxed{f'}} \cdot \overbrace{\sin x}^{\boxed{g}} dx = x \sin x + \cos x + C$$

Exercise

Exercise 9-2 Evaluate the following.

$$(1) \int \cos^2 x \sin x dx \quad (2) \int x e^{6x} dx$$

Ans.

Pause the video and solve by yourself.

Answers to the Exercise

Exercise 9-2 Evaluate the followings.

$$(1) \int \cos^2 x \sin x dx \quad (2) \int x e^{6x} dx$$

Ans. (1) Since $(\cos x)' = -\sin x$, we put $\cos x = u$

$$\therefore -\sin x = \frac{du}{dx}$$

$$\text{Therefore } \int \cos^2 x \sin x dx = \int u^2 (-du) = -\frac{1}{3}u^3 + C = -\frac{1}{3}\cos^3 x + C$$

$$(2) \int \underbrace{x}_{f} \underbrace{e^{6x}}_{g'} dx = x \left(\frac{e^{6x}}{6} \right) - \int 1 \cdot \left(\frac{e^{6x}}{6} \right) dx = \frac{x}{6} e^{6x} - \frac{x}{36} e^{6x} + C$$