

Lesson 11

Estimating Area by Rectangles


11A

- Estimating Area by Rectangles



Estimation of Area by Rectangles

Calculate the area A between $y = x^2 + 1$ and $y = 0$ on $[0, 2]$.
(Suppose that we don't know the definite integral.)

Spritz $[0, 2]$ into 4 subintervals.



$$\begin{aligned} A_R &= \frac{1}{2}f\left(\frac{1}{2}\right) + \frac{1}{2}f(1) + \frac{1}{2}f\left(\frac{3}{2}\right) + \frac{1}{2}f(2) \\ &= \frac{1}{2}\left(\frac{5}{4}\right) + \frac{1}{2}(2) + \frac{1}{2}\left(\frac{13}{4}\right) + \frac{1}{2}(5) = 5.75 \end{aligned}$$




$$\begin{aligned} A_L &= \frac{1}{2}f(0) + \frac{1}{2}f\left(\frac{1}{2}\right) + \frac{1}{2}f(1) + \frac{1}{2}f\left(\frac{3}{2}\right) \\ &= \frac{1}{2}(1) + \frac{1}{2}\left(\frac{5}{4}\right) + \frac{1}{2}(2) + \frac{1}{2}\left(\frac{13}{4}\right) = 3.75 \end{aligned}$$

$$3.75 < A < 5.75$$

Estimation of Area by Rectangles – Cont.

Sprit $[0, 2]$ into 8 subintervals.



$$4.1875 < A < 5.1875$$



Exact value $A = \int_0^2 (x^2 + 1) dt = \left[\frac{x^3}{3} + x \right]_0^2 = \frac{14}{3} = 4.666$

Theorem


If $f(x)$ is continuous on $[a, b]$ then the right and left endpoints approximations approach one and the same limit as $N \rightarrow \infty$

$$\lim_{N \rightarrow \infty} A_R = \lim_{N \rightarrow \infty} A_L = L$$


Example

Examples 11-1 Find the area between $y = x^2$ and $y = 0$ on $[0, 1]$ by making $n \rightarrow \infty$. Use the following formula.

$$1^2 + 2^2 + 3^2 + \dots + n^2 = \frac{n(n+1)(n+2)}{6}$$



Ans.




$$\begin{aligned} S_n &= \frac{1}{n} \left\{ 0 + \left(\frac{1}{n}\right)^2 + \left(\frac{2}{n}\right)^2 + \dots + \left(\frac{n+1}{n}\right)^2 \right\} \\ &= \frac{1}{n^3} \{ 1^2 + 2^2 + \dots + (n-1)^2 \} \\ &= \frac{1}{n^3} \cdot \frac{1}{6} (n-1)n(2n-1) \end{aligned}$$

Therefore

$$\lim_{n \rightarrow \infty} S_n = \frac{1}{3}$$

<Compare>
$$S = \int_0^1 x^2 dt = \left[\frac{x^3}{3} \right]_0^1 = \frac{1}{3}$$

Historical Definition of the Definite Integral



$$a = x_0, x_1, x_2, \dots, x_n = b$$

$$\text{Width } \Delta x = (b - a) / n$$

The right-endpoint approximation:

$$A_{Rn} = f(x_1)\Delta x + f(x_2)\Delta x + \dots + f(x_n)\Delta x$$

$$= \sum_{k=1}^n f(x_k) \Delta x$$

(**Riemann sum** of f over $[a, b]$).

Historical definition of the definite integral

Left-endpoint approx.

$$\int_a^b f(x) dx = \lim_{n \rightarrow \infty} \sum_{k=0}^{n-1} f(x_k) \Delta x$$

Right-endpoint approx.

$$\int_a^b f(x) dx = \lim_{n \rightarrow \infty} \sum_{k=1}^n f(x_k) \Delta x$$

Lesson 1 1

Estimating Area by Rectangles

1 1 B

- Finding of the Limit Value of a Series

Finding the Limit Value of a Series by an Integral

If we put $a = 0$, $b = 1$, we have

$$\int_0^1 f(x) dx = \lim_{n \rightarrow \infty} \frac{1}{n} \sum_{k=0}^{n-1} f\left(\frac{k}{n}\right) \quad \text{Left-endpoint approx.}$$
$$= \lim_{n \rightarrow \infty} \frac{1}{n} \sum_{k=1}^n f\left(\frac{k}{n}\right) \quad \text{Right-endpoint approx.}$$

This relationship is sometimes used to find the sum of infinite series.

Example

Examples 11-2 Find the limit value of

$$S = \lim_{n \rightarrow \infty} \left(\frac{1}{n+1} + \frac{1}{n+2} + \frac{1}{n+3} + \dots + \frac{1}{2n} \right)$$


Ans. Take out quantity $\frac{1}{n}$ from all terms.

$$S = \lim_{n \rightarrow \infty} \frac{1}{n} \left(\frac{1}{1+\frac{1}{n}} + \frac{1}{1+\frac{2}{n}} + \frac{1}{1+\frac{3}{n}} + \dots + \frac{1}{1+\frac{n}{n}} \right)$$

$$= \lim_{n \rightarrow \infty} \frac{1}{n} \sum_{k=1}^n \left(\frac{1}{1+\frac{k}{n}} \right)$$

Replace the part $\frac{k}{n}$ by x and make the definite integral

$$S = \int_0^1 \frac{dx}{1+x} = \left[\log(1+x) \right]_0^1 = \log 2$$



Ex.11-1

Find the value of the following series.

$$S = \lim_{n \rightarrow \infty} \frac{1}{n} \left(\sin \frac{\pi}{n} + \sin \frac{2\pi}{n} + \sin \frac{3\pi}{n} + \dots + \sin \frac{n\pi}{n} \right)$$

Ans.

Pause the video and solve the problem by yourself.

Ex.11-1

Find the value of

$$S = \lim_{n \rightarrow \infty} \frac{1}{n} \left(\sin \frac{\pi}{n} + \sin \frac{2\pi}{n} + \sin \frac{3\pi}{n} + \dots + \sin \frac{n\pi}{n} \right)$$

Ans.

$$\begin{aligned} S &= \lim_{h \rightarrow \infty} \frac{1}{n} \sum_{k=1}^n \sin \left(\frac{k}{n} \pi \right) = \int_0^1 \sin(\pi x) dx = \left[-\frac{\cos(\pi x)}{\pi} \right]_0^1 \\ &= -\left(\frac{\cos \pi - 1}{\pi} \right) = \frac{2}{\pi} \end{aligned}$$